

References and Notes

1. F. C. Fuglister, in *Studies in Physical Oceanography. A Tribute to Georg Wüst on His 80th Birthday*, A. L. Gordon, Ed. (Gordon & Breach, New York, 1972), vol. 1, p. 137.
2. C. E. Parker, *Deep-Sea Res.* **18**, 981 (1971).
3. D. Y. Lai and P. L. Richardson, *J. Phys. Oceanogr.* **7**, 670 (1977).
4. P. L. Richardson, R. E. Cheney, L. V. Worthington, *J. Geophys. Res.* **83**, 6136 (1978).
5. H. Kawai, in *Proceedings of the Fourth Cooperative Study, Kuroshio (CSK), Tokyo, February 1979* (Saikon, Tokyo, 1979), p. 250.
6. C. S. Nilsson, J. C. Andrews, P. Scully-Power, *J. Phys. Oceanogr.* **7**, 659 (1977).
7. R. E. Cheney, W. H. Gemmill, M. K. Shank, P. L. Richardson, D. Webb, *ibid.* **6**, 741 (1976).
8. P. H. Wiebe, K. H. Burt, S. H. Boyd, A. W. Morton, *J. Mar. Res.* **34**, 313 (1976).
9. R. A. Dohlar and R. E. Cheney, *J. Phys. Oceanogr.* **7**, 944 (1977).
10. A. Leetmaa, *Science* **198**, 188 (1977).
11. A convenient way of expressing the magnitude of the cold-core ring anomaly is to indicate the minimum depth to 15°C at the ring center. The Slope Water-Gulf Stream boundary is commonly determined by the 15°C isotherm at 200 m, with this isotherm being shallower to the north and deeper to the south; thus, the shallower the 15°C isotherm in a cold-core ring the greater the anomaly.
12. N. E. Huang, C. D. Leita, C. G. Parra, *J. Geophys. Res.* **83**, 4673 (1978).
13. R. E. Cheney and J. G. Marsh, in preparation.
14. D. E. Hagan, D. B. Olson, J. E. Schmitz, A. C. Vastano, *J. Phys. Oceanogr.* **8**, 997 (1978).
15. P. Mukherji and D. R. Kester, *Science* **204**, 64 (1979).
16. F. A. Richards and A. C. Redfield, *Deep-Sea Res.* **2**, 182 (1955).
17. We use here the same terminology as in Backus *et al.* (43).
18. P. L. Richardson, *J. Phys. Oceanogr.* **10**, 90 (1980).
19. C. Maillard, T. B. Sanford, *J. Geophys. Res.* **84**, 7727 (1979).
20. F. C. Fuglister, in *A Voyage of Discovery, George Deacon 70th Anniversary Volume*, M. V. Angel, Ed. (Pergamon, New York, 1977), p. 177.
21. A. C. Vastano, J. E. Schmitz, D. E. Hagan, *J. Phys. Oceanogr.* **10**, 493 (1980).
22. C.-G. Rossby, *J. Mar. Res.* **2**, 38 (1939).
23. B. A. Warren, *Deep-Sea Res.* **14**, 505 (1967).
24. J. McWilliams and G. R. Flierl, *J. Phys. Oceanogr.* **9**, 1155 (1979).
25. R. Mied and G. J. Lindemann, *ibid.*, p. 1183.
26. A. C. Vastano, D. E. Hagan, J. E. Schmitz, in preparation.
27. J. R. Barrett, *Deep-Sea Res.* **18**, 1221 (1971).
28. R. E. Cheney and P. L. Richardson, *ibid.* **23**, 143 (1976).
29. G. R. Flierl, *J. Phys. Oceanogr.* **7**, 365 (1977).
30. D. B. Olson, *ibid.* **10**, 514 (1980).
31. D. C. Smith, thesis, Texas A & M University (1980).
32. R. B. Lambert, Jr., *Deep-Sea Res.* **21**, 529 (1974).
33. R. L. Molinari, thesis, Texas A & M University (1970).
34. J. E. Schmitz and A. C. Vastano, *J. Phys. Oceanogr.* **5**, 93 (1975).
35. A. C. Vastano and D. E. Hagan, *ibid.* **7**, 938 (1977).
36. P. H. Wiebe and S. H. Boyd, *J. Mar. Res.* **36**, 119 (1978).
37. A. Fleminger and K. Hulsemann, *Mar. Biol.* **40**, 233 (1977).
38. P. H. Wiebe, E. M. Hulbert, E. J. Carpenter, A. E. Jahn, G. P. Knapp III, S. H. Boyd, P. B. Ortner, J. L. Cox, *Deep-Sea Res.* **23**, 695 (1976).
39. P. B. Ortner, P. H. Wiebe, L. R. Hauray, S. H. Boyd, *Fish. Bull.* **76**, 323 (1978).
40. S. H. Boyd, P. H. Wiebe, J. L. Cox, *J. Mar. Res.* **36**, 143 (1978).
41. P. B. Ortner, E. M. Hulbert, P. H. Wiebe, *J. Exp. Mar. Biol. Ecol.* **39**, 101 (1979).
42. P. B. Ortner, P. H. Wiebe, J. L. Cox, *J. Mar. Res.* **38**, 507 (1980).
43. R. H. Backus, J. E. Craddock, R. L. Haedrich, B. H. Robison, C. E. Karnella, *Mem. Sears Found. Mar. Res.* **1** (part 7), 266 (1977).
44. B. G. Nafpaktitis, R. H. Backus, J. E. Craddock, R. L. Haedrich, B. H. Robison, C. E. Karnella, *ibid.*, p. 38.
45. C. Karnella, personal communication.
46. D. R. Watts and D. B. Olson, *Science* **202**, 971 (1978).
47. W. R. Holland, *J. Phys. Oceanogr.* **8**, 363 (1978).
48. In accordance with Lai and Richardson (3), we assume that the area affected by cold-core rings is that part of the northern Sargasso Sea west of 50°W, about 3×10^{12} m², and the volume affected is that part of the area above the permanent thermocline, or about 3×10^{15} m³. The southern boundary of the northern Sargasso Sea is judged to lie at about 28°N, corresponding to the southern limit of the principal westward return flow of the Gulf Stream (49).
49. L. V. Worthington, *Johns Hopkins Oceanogr. Stud.* **6** (1976), figure 42.
50. C. Wunsch, *Rev. Geophys. Space Phys.* **16**, 583 (1978).
51. W. J. Schmitz, Jr., *J. Mar. Res.* **38**, 111 (1980).
52. G. T. Rowe and W. D. Gardner, *ibid.* **37**, 581 (1979).
53. R. G. Fairbanks, P. H. Wiebe, A. W. H. Bé, *Science* **207**, 61 (1980).
54. MODE group, *Deep-Sea Res.* **25**, 859 (1978).
55. P. H. Wiebe, S. H. Boyd, J. L. Cox, *Fish. Bull.* **73**, 777 (1975).
56. This work was largely supported through contracts with the Office of Naval Research. Additional funding was provided by the National Science Foundation. We especially would like to thank our shipmates on the Research Vessels *Knorr* and *Endeavor*, the GEOSECS Operations Group at the Scripps Institution of Oceanography, especially the late Arnold Bainbridge, for help in making the CTD profiles, and also the following, who assisted with data collection, processing, and analysis: Steven H. Boyd, Robert H. Byrne, James L. Cox, James E. Craddock, Denise E. Hagan, G. Lee Johnson, Charles E. Lea, Alfred W. Morton, P. Mukherji, Joyce E. Schmitz, and Richard W. Zuehlke. The present address of D.B.D. is: Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Fla. 33149.

Psychoneuroendocrine Influences on Immunocompetence and Neoplasia

Vernon Riley

"Stress" is a widely used term for describing emotional and biological responses to novel or threatening situations. There is, thus, an extensive variety of experimental or other circumstances in which "stress" serves as a convenient word to express complex and incompletely understood psychological and physiological phenomena (1-3).

In studies at this laboratory, we use the term "stress" in a more restricted experimental sense to relate specific

stress-inducing stimuli, or stressors, to their physiological consequences. The latter include specific biochemical, cellular, and tissue alterations that are associated with an emotional activation of the adrenal cortex by way of the pituitary and its secretion of adrenocorticotropic hormone (4, 5). Within the biological systems that we have used, several key parameters characterize the physiological manifestations of stress, and relate to pathological and other changes that may be observed in stressed experimental an-

imals measured with precision the most conspicuous, and what appears to be the most relevant, of the biochemical substances elaborated by this organ in response to anxiety, namely, corticosterone. Immediately after an animal is subjected to an emotional stimulus, or perceives a situation that generates anxiety, the adrenal cortex in response to signals from the hypothalamus, via the pituitary, produces increased quantities of corticosterone. The rapidity of the appearance of corticosterone in the plasma can be readily measured by appropriate microassay techniques (6-8).

Immunological and Pathological Consequences of Stress

Secondary manifestations that result from increased corticosterone in the blood plasma that are readily observed include (i) lymphocytopenia, or decreased circulating lymphocytes, (ii) thymus involution, and (iii) related loss of tissue mass of the spleen and peripheral