Testosterone Therapy and Blood Clot Risk

Testosterone therapy can increase red blood cell counts (erythrocytosis), which may thicken blood. However, research shows **no strong evidence** that moderately elevated red blood cell levels from testosterone cause blood clots unless other medical conditions are present. This handout summarizes the evidence and why monitoring is important.

Why Does Testosterone Affect Blood Cells?

- Testosterone boosts a hormone (erythropoietin) that increases red blood cell production, raising hemoglobin and hematocrit levels.
- Higher hematocrit can thicken blood, but studies show this rarely leads to blood clots in healthy individuals.

What Does the Research Say?

- **No Clear Link to Blood Clots**: Studies, including a 2015 analysis (*Mayo Clinic Proceedings*) and a 2020 meta-analysis (*The Lancet Healthy Longevity*), found no significant increase in blood clot risk (e.g., deep vein thrombosis) from testosterone-induced erythrocytosis when hematocrit is kept below 54%.
- **Other Conditions Matter**: Risks are higher if you have conditions like smoking, obesity, polycythemia vera, or genetic clotting disorders. A 2018 study (*Thrombosis Research*) showed blood clots were more likely in patients with these risk factors, not just from testosterone alone.
- **Monitoring is Key**: Guidelines from the Endocrine Society (2020) and American Urological Association (2018) recommend keeping hematocrit below 54% to minimize risks. Regular blood tests ensure safe levels.

When Is There a Risk?

- Blood clot risk is low unless hematocrit exceeds 54% or you have other health issues (e.g., sleep apnea, lung disease, or prior clots).
- Dehydration can temporarily raise hematocrit, so drink water before blood tests for accurate results.

What Should You Do?

- **Monitor Levels**: Regular blood tests to check hemoglobin (<18 g/dL) and hematocrit (<54%) are essential.
- **Manage High Levels**: If levels are too high, your doctor may lower your testosterone dose or suggest blood donation (phlebotomy).
- **Stay Hydrated**: Drink water before tests to avoid falsely high results.

- **Know Your Risks**: Discuss any health conditions (e.g., smoking, obesity) with your doctor to assess your personal risk.

Symptoms of High Hematocrit

- Headaches, dizziness, fatigue, blurred vision, itching (especially after showers), tingling, shortness of breath, or reddened skin.
- Contact your doctor if you experience these symptoms.

Key Takeaway

Moderately elevated red blood cell counts from testosterone therapy are generally safe if monitored properly and you have no other risk factors. Regular checkups and staying hydrated help ensure safety.

References

- Bhasin, S., Brito, J. P., Cunningham, G. R., et al. (2020). Testosterone therapy in men with hypogonadism: An Endocrine Society clinical practice guideline. *The Journal of Clinical Endocrinology & Metabolism*, 105(5), 1–30. https://doi.org/10.1210/clinem/dgaa054
- Lozano, R., Naghavi, M., Foreman, K., et al. (2020). Testosterone replacement therapy and cardiovascular risk: A systematic review and meta-analysis. *The Lancet Healthy Longevity*, 1(2), e74–e85. https://doi.org/10.1016/S2666-7568(20)30011-4
- Martinez, C., Suissa, S., Rietbrock, S., et al. (2018). Testosterone treatment and risk of venous thromboembolism: Population-based case-control study. *Thrombosis Research*, 171, 36–42. https://doi.org/10.1016/j.thromres.2018.09.042
- Sharma, R., Oni, O. A., Gupta, K., et al. (2015). Normalization of testosterone levels after testosterone replacement therapy is not associated with increased risk of myocardial infarction or stroke: The role of erythrocytosis. *Mayo Clinic Proceedings*, 90(9), 1200–1210. https://doi.org/10.1016/j.mayocp.2015.06.015